

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 7/00	A1	(11) International Publication Number: WO 95/23578 (43) International Publication Date: 8 September 1995 (08.09.95)
(21) International Application Number: PCT/EP95/00716 (22) International Filing Date: 25 February 1995 (25.02.95) (30) Priority Data: 94301574.3 4 March 1994 (04.03.94) EP <i>(34) Countries for which the regional or international application was filed:</i> GB et al. (71) Applicant (for AU BB CA GB IE KE LK MN MW NZ SD SZ TT UG only): UNILEVER PLC (GB/GB); Unilever House, Blackfriars, London EC4 4BQ (GB). (71) Applicant (for all designated States except AU BB CA GR IE KE LK MN MW NZ SD SZ TT UG): UNILEVER NV (NL/NL); Weena 455, NL-3013 AL Rotterdam (NL). (72) Inventors: FINEL, Christophe, Michel; 221 Bebington Road, Rock-Ferry, Wirral, Merseyside L42 2QA (GB). HAGUE, Jonathan, David; 24 Trafalgar Drive, Bebington, Wirral, Merseyside L63 7RW (GB). REID, Euan, Stuart; 4 Meadway, Bebington, Wirral, Merseyside L62 2 AR (GB). (74) Agent: MULDER, C., W., R.; Unilever plc. Patent Division, Colworth House, Sharnbrook, Bedford. MK44 1LQ (GB).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TT, UA, UG, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). ARIPO patent (KE, MW, SD, SZ, UG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: LIPOSOMES FOR DEPOSITION ON HAIR		
(57) Abstract		
Process for the deposition, of an active ingredient on the hair by cationic liposomes, and post-wash conditioner or hair treatment masque compositions containing cationic liposomes and an active ingredient.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauntania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

LIPOSOMES FOR DEPOSITION ON HAIR

FIELD OF THE INVENTION

5 This invention concerns processes and compositions for the treatment of human hair. More particularly the invention concerns an improved system for the deposition of active ingredients on hair from hair treatment compositions.

10 BACKGROUND OF THE INVENTION AND PRIOR ART

When treating hair with rinse-off products incorporating surfactant soluble active ingredients, a considerable amount of the active ingredient will be rinsed away.

15 Studies have shown that the level of retention in the case of a simple shampoo composition, where the active ingredient is solubilised in the surfactant micelles in the product, can be as low as 5%. Retention of these active ingredients from conditioning compositions is generally
20 higher, probably because there is poorer solubilisation of the active ingredient in the conditioning base, but there is considerable scope for improvement, as this would provide better performance of the active ingredient and the option of reducing the level of expensive active ingredient
25 in the product, with consequent cost saving.

It is an object of the present invention to provide an improved system for the deposition of active ingredients on hair.

30

BRIEF SUMMARY OF THE INVENTION

We have now found that cationic liposomes, ie. aqueous compartments enclosed by one or more lipid bilayers, can be
35 formed which are storage-stable and possess an affinity for hair. When included in a hair treatment composition, these

- 2 -

liposomes deposit on the hair, are not eliminated during rinsing and can enhance the deposition on the hair of active ingredients with which they are combined.

5 Accordingly, in a first aspect, the present invention provides a process for the deposition of an active ingredient on hair by cationic liposomes.

10 In a second aspect, the present invention provides a composition comprising cationic liposomes and an active ingredient, which when applied to hair will cause enhanced deposition of the active ingredient on the hair than previously achievable.

15 DETAILED DESCRIPTION OF THE INVENTION

A preferred process according to the invention comprises the following steps:

20 (a) forming a dispersion of cationic liposomes incorporating the active ingredient,

(b) processing the dispersion into a hair treatment composition, and

25 (c) treating the hair with the composition.

We have found that the dispersion of step(a) may be prepared by the simple addition of solid cholesterol ,
30 along with the active ingredient, to an aqueous solution of cationic surfactant. This induces the formation of liposomes, and the structures so generated may be easily visualised using conventional contrast microscopy techniques.

35

- 3 -

The concentration of cationic surfactant in the liposomal dispersion is suitably from 2% to 10% by weight based on total weight, and the weight ratio of cholesterol to cationic surfactant is preferably 1:1.

5

Microscopy studies of liposomal dispersions prepared as above demonstrate at least partial encapsulation of the active ingredient in the liposomes. It is preferable to optimise such encapsulation to allow the best efficiency of deposition enhancement from the liposomal dispersions of the invention, although the exact mechanism by which this occurs is unclear. In general, it is less preferable to add the active ingredient at a later stage than addition of the liposomes, since this appears to be detrimental to encapsulation of the active ingredient by the liposomes.

10
15

However, in an alternative process according to the invention, a dispersion of cationic liposomes may be formed in situ in a hair treatment composition by the inclusion of liposomal components, typically cholesterol and cationic surfactant, during the processing stage. The active ingredient may then be incorporated into the composition after this stage.

20

In this alternative process, the concentration of cholesterol added during processing (by weight based on the total weight of the hair treatment composition) is suitably from 0.05 to 3%. e.g. 0.1 to 1% and the concentration of cationic surfactant added during processing is suitably from 0.15 to 5%, e.g. 0.5 to 2%, by weight based on total weight of the hair treatment composition.

25
30

Surprisingly, this also provides a favourable route to effective liposomal encapsulation of the active ingredient. It appears that in such a case, the whole of the compositional microstructure is altered so as to represent

35

- 4 -

a favourable environment for active ingredient encapsulation.

The active ingredient is normally a water-insoluble
5 or sparingly water-soluble substance, such as an oil which may take the form of a sunscreen. Among suitable sunscreens are, camphor derivatives, benzophenone compounds such as 4,4'-tetrahydroxy-benzophenone, sold commercially as Uvinul D50, and 2-hydroxy-4-methoxybenzophenone, sold commercially
10 as Eusolex 4360, dibenzoyl methane derivatives such as t-butyl-4-methoxydibenzoylmethane, sold commercially as Parsol 1789, and isopropylidibenzoyl methane, sold commercially as Eusolex 8020. Preferred sunscreen materials are cinnamates, such as 2-ethylhexyl-p-methoxy cinnamate,
15 sold commercially as Parsol MCX, 2-ethoxy ethyl-p-methoxy cinnamate, sold commercially as Giv-Tan F and isoamyl-p-methoxy cinnamate, sold commercially as Neo-Heliopan E1000.

The active ingredient may also be an antidandruff agent,
20 such as zinc pyrithione, and other 1-hydroxy pyridones. A preferred antidandruff agent is the 1-hydroxy-2-pyridone derivative known as piroctone olamine, whose chemical name is 1-hydroxy-4-methyl-6-(2,4,4-trimethylpentyl)-2-pyridone monoethanolamine salt, and which is sold under the trade
25 name OCTIPIROX by Hoechst AG.

Other suitable active ingredients are vitamin E and derivatives thereof, volatile oils and perfumes.

30 The active ingredient is normally present in a concentration of from 0.005% to 5%, preferably from 0.085% to 2% by weight based on the total weight of the hair treatment composition. The optimum concentration of active
35 ingredient will depend on the precise chemical nature of the active ingredient. For a sunscreen, if the composition comprises less than 0.005% by weight of the sunscreen,

- 5 -

little benefit will be obtained and if greater than 5% is present, it is unlikely that additional benefit will be obtained.

5 Examples of suitable cationic surfactants include:
quaternary ammonium hydroxides, e.g.
tetramethylammonium hydroxide, alkyltrimethylammonium
hydroxides in which the alkyl group has from about 8 to 22
10 carbon atoms, for example octyltrimethylammonium hydroxide,
dodecyltrimethylammonium hydroxide,
hexadecyltrimethylammonium hydroxide,
cetyltrimethylammonium hydroxide and
behenyltrimethylammonium hydroxide,
benzyltrimethylammonium hydroxide,
15 octyldimethylbenzylammonium hydroxide,
decyldimethylbenzylammonium hydroxide,
stearyldimethylbenzylammonium hydroxide,
didodecyldimethylammonium hydroxide,
dioctadecyldimethylammonium hydroxide, tallow
20 trimethylammonium hydroxide, cocotrimethylammonium
hydroxide, and their corresponding salts, e.g. halides
Cetylpyridinium hydroxide or its corresponding salts, e.g.
halide.

25 Preferred cationic surfactants are cetyl trimethylammonium
chloride and cetyltrimethylammonium bromide, hereinafter
referred to as C.T.A.C. and C.T.A.B. respectively.

30 Compositions according to the present invention may further
comprise one or more optional ingredients which are
normally found in hair treatment compositions. The
compositions of the invention will preferably take the form
of post-wash hair conditioning compositions or hair
treatment masques, but may also take the form of
35 conditioning shampoos or hair styling compositions or the
like.

- 6 -

One preferred optional component which may be included in the hair treatment compositions of the invention is a fatty alcohol or fatty acid, or derivative thereof, or a mixture of any of these, having a chain length of from about 8 to
5 about 28 carbon atoms, more preferably from about 12 to about 18 carbon atoms. These materials may be predominantly linear or may be branched.

Such fatty material(s) may be present in the compositions
10 of the invention in a total amount of from about 0.001 to 20% by weight, more preferably 0.01 to 10%, even more preferably 0.01 to 5% yet more preferably 0.1 to 1%. An especially preferred amount of the fatty material, if
15 present, is up to about 0.5% by weight, since such amounts help to render the compositions smooth textured and non-lumpy.

Where it is desired to formulate a hair treatment composition of the invention which not only has
20 conditioning properties but also has detergent properties, i.e. a shampoo, then one or more surfactants may be included, preferably selected from nonionic, amphoteric and zwitterionic surfactants.

25 Nonionic surfactants suitable for use in compositions of the invention include condensation products of aliphatic C_6-C_{18} primary or secondary linear or branched chain alcohols or phenols with alkylene oxides, usually ethylene oxide, and generally having from 6 to 30 ethylene oxide
30 groups.


Other suitable nonionics include mono- or di-alkyl alkanolamides. Examples include coco mono- or di-ethanolamide and coco mono-isopropanolamide.

35

- 7 -

Further suitable nonionic surfactants are the alkyl polyglycosides (APG's). Typically, the APG is one which comprises an alkyl group connected (optionally via a bridging group) to a block of one or more glycosyl groups.

5 Preferred APG's are described by the following formula:



10

Wherein R is a branched or straight chain alkyl group which may be saturated or unsaturated and G is a saccharide group.

15

R may represent a mean alkyl chain length of from about C_5 to about C_{20} . Preferably, R represents a mean alkyl chain length of from about C_8 to about C_{12} . Most preferably the value of R lies between about 9.5 and about 10.5. G may be selected from C_5 or C_6 monosaccharide residues or mixtures

20 of C_5 and C_6 monosaccharide residues, and is preferably a glucoside. G may be selected from the group comprising glucose, xylose, lactose, fructose, mannose and derivatives thereof. Preferably G is glucose.

25

The degree of polymerisation, n, may have a value of from about 1 to about 10 or more. Preferably, the value of n lies in the range of from about 1.1 to about 2. Most preferably the value of n lies in the range of from about 1.3 to about 1.5.

30

Suitable alkyl polyglycosides for use in the invention are commercially available and include for example those materials identified as: Oramix NS10 ex Seppic; and APG225, APG300, APG350, APG550 and APG600 ex Henkel.

35

- 8 -

Amphoteric and zwitterionic surfactants suitable for use in compositions of the invention may include alkyl amine oxides, alkyl betaines, alkyl amidopropyl betaines, alkyl sulphobetaines (sultaines), alkyl glycinate, alkyl carboxyglycinates, alkyl amphopropionates, alkylamphoglycinates alkyl amidopropyl hydroxysultaines, acyl taurates and acyl glutamates, wherein the alkyl and acyl groups have from 8 to 19 carbon atoms. Examples include lauryl amine oxide, cocodimethyl sulphopropyl betaine and preferably lauryl betaine, cocamidopropyl betaine and sodium cocamphopropionate.

Further surfactants which may be suitable for use in shampoos in accordance with the invention include one or more anionic surfactants instead of or in addition to any of those surfactants mentioned above.

Suitable anionic surfactants are the alkyl sulphates, alkyl ether sulphates, alkaryl sulphonates, alkanoyl isethionates, alkyl succinates, alkyl sulphosuccinates, N-alkoyl sarcosinates, alkyl phosphates, alkyl ether phosphates, alkyl ether carboxylates, and alpha-olefin sulphonates, especially their sodium, magnesium, ammonium and mono-, di- and triethanolamine salts. The alkyl and acyl groups generally contain from 8 to 18 carbon atoms and may be unsaturated. The alkyl ether sulphates, alkyl ether phosphates and alkyl ether carboxylates may contain from one to 10 ethylene oxide or propylene oxide units per molecule, and preferably contain 2 to 3 ethylene oxide units per molecule.

Examples of suitable anionic surfactants include sodium oleyl succinate, ammonium lauryl sulphosuccinate, ammonium lauryl sulphate, sodium dodecylbenzene sulphonate, triethanolamine dodecylbenzene sulphonate, sodium cocoyl isethionate, sodium lauroyl isethionate and sodium N-lauryl

- 9 -

sarcosinate. The most preferred anionic surfactants are sodium lauryl sulphate, triethanolamine lauryl sulphate, triethanolamine monolauryl phosphate, sodium lauryl ether sulphate 1EO, 2EO and 3EO, ammonium lauryl sulphate and ammonium lauryl ether sulphate 1EO, 2EO and 3EO.

The surfactant(s) may be present in the hair treatment composition in a total amount of from about 1 to 70% by weight, preferably from 2 to 40% by weight, more preferably from 5 to 30% by weight.

As further optional components for inclusion in the compositions of the invention, in addition to water, the following may be mentioned: pH adjusting agents, viscosity modifiers, pearlescers, opacifiers, suspending agents, preservatives, colouring agents, dyes, proteins, herb and plant extracts, polyols and other moisturising and/or conditioning agents.

Embodiments of the present invention will now be further illustrated by reference to the following examples. All amounts given are in % by weight, unless otherwise stated.

EXAMPLES:

25

Materials

Cholesterol was 95%, as supplied by Aldrich Chemical Co. CTAB was 99%, as supplied by Fluka AG. CTAC was obtained as a 50% solution (Arquad 16-50, obtained from AKZO). Octipirox and Parsol MCX were obtained from Hoescht and Givaudan-Roure respectively.

- 10 -

MethodsLiposome Preparation

5 CTAC was added to water at room temperature. Solid
cholesterol was added, along with either Octipirox or
Parsol MCX. The mixture was dispersed for three minutes
with high shear at room temperature using a Silverson
mixer. Liposome preparations were always 1:1 CTAC:
10 Cholesterol by weight, and were between 2 and 10% CTAC.

Examples 1 to 3Investigation of the affinity of liposomes for hair

15 A liposome solution was prepared as above using the
following ingredients:

Cholesterol 0.125% (estimated total liposome concentration:
20 0.25%)
CTAC 0.125%
Demineralized water

Procedure

25 Hair switches of the same quality weighing about 1g were
selected, and each switch immersed separately in 15ml of
the prepared liposome solution. After removal of the
switch, the concentration of each solution was investigated
30 by measuring UV absorbance at 400nm. The system was
calibrated with two solutions having liposome concentration
of 0.20% and 0.25%.

Results are given in Table 1:

35

- 11 -

Table 1

Example	Switch immersion time	Final concentration of the solution
1	15 minutes	0.246%
2	2 hours	0.233%
3	16 hours	0.231%

Conclusion

If all of the components in the solution are assumed to be in the form of liposomes, the results indicate that liposomes are deposited on hair, as a result of the affinity between the liposomes and the hair.

Examples 4 and 5

Parsol MCX uptake by hair was examined:

- from the liposomal dispersion alone, and from micellar CTAB solution.

Procedure

Hair switches were treated with 0.2g conditioner/g hair, and 4 x 4g switches were employed. Switches were extracted after rinsing with 100 ml of ethanol for 1 hour.

Parsol concentration in the ethanol solutions were determined using a calibration curve generated by several dilutions (in ethanol) of the initial liposome dispersion, or of Parsol MCX in ethanol. Absorbances were measured at a wavelength of 309 nm.

- 12 -

Table 2 displays retention results obtained from:

Comparative Example A : 4% CTAB, 0.3% Parsol MCX

Example 4 : 4% CTAB, 4% Cholesterol, 0.3%
Parsol MCX.

Comparative Example B : 2% CTAB, 0.15% Parsol MCX

Example 5 : 2% CTAB, 2% Cholesterol, 0.15%
Parsol MCX

In Comparative Examples A and B, the Parsol MCX was shown
by microscopy to be fully solubilised in the surfactant
micelles. In Examples 4 and 5, the Parsol MCX was judged
from microstructural studies to have been completely
incorporated into the liposomal dispersion.

Table 2

Example	Parsol Retained (ug/g Hair)	Efficiency of Parsol Retention
4	95	15.7
Comparative Example A	75	12.7
5	55	17.9
Comparative Example B	45	14.4

For each of the CTAB concentrations tested, addition of
cholesterol, which induces liposomes in the system, allows
a 25% increase in Parsol MCX retention.

- 13 -

Examples 6 to 8

Parsol MCX uptake was examined:

- 5 - from fully formulated conditioners* with 10% post-processing additions of :

Comparative Example C : 2% Parsol MCX in distilled water,

10 Example 6 : 7.5% CTAC, 7.5% Cholesterol, 2% Parsol MCX

 Example 7 : 10% CTAC, 10% Cholesterol, 2% Parsol MCX,

and

15 Example 8 : 4% CTAC, 4% Cholesterol, 0.85% Parsol

* The conditioner base to which the post-processing additions were made was made up from the following ingredients:

20

	Ingredient	%wt
	CTAC	0.7%
	Cetearyl alcohol	3.5%
	paraffin	1%
25	glyceryl stearate	0.7%
	butyl hydroxy toluene (BHT)	0.05%
	Bronopol (2-bromo-nitropropane-1,3-diol)	0.01%
	perfume	0.2%
	silk amino acids	0.2%
30	demineralised water	83.44%

- 14 -

Procedure

Hair switches (1g) were treated with 1g of conditioner + post-processing additive for 3 minutes, followed by a rinse of 1 minute. Switches were then extracted for 30 seconds with 10 ml of ethanol. Ethanolic extractions from the treated hair were diluted x15 with ethanol prior to measurement of absorbance.

Retention results are displayed in Table 3.

Table 3

Example	Parsol Retained (ug/g Hair)	Efficiency of Parsol Retention (%)
Comparative Example C	800	4.1
Example 6	1,150	5.6
Example 7	1,550	7.7
Example 8	800	9.6

The results demonstrate the advantage of processing Parsol MCX into the dispersion of CTAC/Cholesterol liposomes. Choosing a higher level of CTAC and cholesterol (Example 7) clearly allows a higher level of Parsol to be retained if the final concentration is 0.2% Parsol MCX. By solubilising a lower level of Parsol MCX (0.85%, giving a final concentration of 0.085%) into 4% CTAC, 4% Cholesterol, (Example 8), a retention level equivalent to that found in the control conditioner containing 0.2% Parsol MCX (Comparative Example C) can be achieved.

- 15 -

Examples 9 to 10**Effect of Method of Liposome Addition on Parsol MCX Uptake.**

5

Parsol MCX retention was studied from the following systems (using a conditioner base made up as described under Examples 6 to 8):

10 **Comparative Example D :** Conditioner with a post-addition of 0.085% Parsol MCX.

Example 9 : Conditioner with a 10% post-addition of a dispersion of 0.85% Parsol MCX, 4% Cholesterol, 4% CTAC.

15 **Comparative Example E :** Conditioner with a 10% post-addition of 4% Cholesterol, 4% CTAC, and with 0.085% Parsol MCX added separately.

20 **Comparative Example F :** Conditioner with 1.1% CTAC (extra) added in the fatty phase during processing, with 0.085% Parsol and 0.4% Cholesterol added post-processing.

25 **Example 10 :** Conditioner with 1.1% CTAC, 0.4% Cholesterol added in the fatty phase during processing, and 0.2% Parsol MCX added post-processing.

30 **Comparative Example G :** Conditioner with 1.1% CTAC added in the fatty phase during processing, and 0.2% Parsol added post-processing. (No cholesterol).

The procedure followed was equivalent to that described above for Examples 6 to 8.

35

Results are detailed in Table 4:

Table 4

Example	Parsol Retained (mg/g hair)	Efficiency of Parsol Retention (%)
Comparative Example D	375	4.1
Example 9	800	9.1
Comparative Example E	625	7.5
Comparative Example F	400	5.2
Example 10	1,825	9.1
Comparative Example G	1,300	6.4

The results suggest that addition of the pre-prepared liposomal components incorporating Parsol MCX after conditioner processing leads to the most effective Parsol MCX retention. However, adding a higher Parsol MCX concentration, after having processed the liposome components into the conditioner (Example 10), is also an effective route to high Parsol MCX retention. In this case, the inclusion of cholesterol at the processing stage seems to have a profound effect on the structure of the conditioner.

- 17 -

Example 11Uptake of Octipirox from fully formulated conditioners

5 Octipirox retention was evaluated from the following conditioning formulations (the conditioner base being made up as described under Examples 6 to 8):

10 Comparative Example H : Conditioner with Octipirox (0.1%) added post-processing in the perfume phase.
Example 11 : Conditioner with a 10% addition of 5% CTAC, 5% Cholesterol and 1% Octipirox.

15 Comparative Example I : Conditioner with a 10% addition of 5% CTAC, 5% Cholesterol, and with 0.1% Octipirox added separately in the perfume phase.

Procedure

20 1g switches were treated with either 1g or 0.2g of conditioner + additive (1 minute application). After a 1 minute rinse, each switch was placed into a centrifuge tube and 10 ml of a solution comprising:

25 4% Acetic acid (12%)
2% Iron (II) sulphate solution (0.17M, acidified with HCl)
94% Methanol

30 was added.

After 35 minutes (+/- 30) seconds in contact with the switch, the mixture was centrifuged at 3500 rpm for 5 minutes. Octipirox content in the supernatant was estimated

- 18 -

from absorbance measurement at 462nm, using a previously determined calibration curve.

The results are shown in Table 5:

5

Table 5

Treatment with 1g conditioner/g hair			Treatment with 0.2g conditioner/g hair		
Example	Parsol Returned (ug/g hair)	Efficiency of Parsol Retention (%)	Example	Parsol retained (ug/g hair)	Efficiency of Parsol Retention (%)
Comparative Example H	850	8.3	Comparative Example H	500	26.3
Example 11	1,800	17.9	Example 11	1,000	46.8
Comparative Example I	1,125	11.4	Comparative Example I	575	27.3

10

15

As was found with Parsol MCX, inclusion of the Octipirox into the liposomal dispersion offers a route to enhancement of retention of x2.

20

Conclusions

25

1) Addition of cholesterol to CTAC or CTAB solutions induces formation of coarse liposomes. The structures generated can easily be visualised using contrast microscopy techniques.

30

2) 1:1 CTAC : cholesterol liposomes are able to partially solubilise an active ingredient such as Parsol MCX or Octipirox.

- 19 -

- 3) When fully incorporated into a liposomal dispersion, Parsol MCX retention on hair can be increased by about 25% relative to retention from a micellar control.
- 5 4) When added as a component of a liposomal premix to a post-wash conditioner, up to 50% enhancement of the retention of both Parsol MCX and Octipirox may be achieved.

10

- 20 -

CLAIMS:

1. A process for the deposition of an active ingredient on hair characterised by applying thereto a hair treatment composition in which the active ingredient is incorporated in cationic liposomes.
2. A process according to claim 1, in which the active ingredient is water-insoluble.
3. A process according to claims 1 or 2 , in which the active ingredient is selected from the group consisting of sunscreens and anti-dandruff agents.
4. A process according to any preceding claim, comprising the following steps:
 - (a) forming a dispersion of cationic liposomes incorporating the active ingredient,
 - (b) processing the dispersion into a hair treatment composition, and
 - (c) treating the hair with the composition.
5. A process according to any of claims 1 to 3, comprising the following steps:
 - (a) forming a dispersion of cationic liposomes in situ in a hair treatment composition by inclusion of liposomal components during processing thereof,
 - (b) incorporating the active ingredient into the composition after processing thereof, and

- 21 -

(c) treating the hair with the composition.

6. A hair treatment composition characterised in that it comprises cationic liposomes and an active ingredient.
- 5 7. A composition according to claim 6 which is formulated as a post-wash conditioner or as a hair treatment masque.
- 10 8. A composition according to claim 6 or claim 7, in which the liposomes are formed from a mixture of cholesterol and cetyltrimethylammonium chloride(C.T.A.C.).
- 15 9. A composition according to claim 8 in which the weight ratio of cholesterol to C.T.A.C. is 1:1.
- 20 10. A composition according to any of claims 6 to 9 in which the active ingredient is selected from the group consisting of Parsol and Octipirox.

INTERNATIONAL SEARCH REPORT

Int. Application No

PCT/EP 95/00716

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K7/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,92 19214 (RICHARDSON-VICKS INC.) 12 November 1992 see the whole document	1-10
X	FR,A,2 676 361 (L'OREAL) 20 November 1992 see the whole document	1-10
X	EP,A,0 373 988 (L'OREAL) 20 June 1990 see the whole document	1-10
A	WO,A,92 08685 (L'OREAL) 29 May 1992 see the whole document	1-10



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

20 July 1995

Date of mailing of the international search report

01.08.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patendaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax (+31-70) 340-3016

Authorized officer

Couckuyt, P

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. .onal Application No

PCT/EP 95/00716

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9219214	12-11-92	US-A- 5229104 AU-A- 1683892 CA-A- 2107076 EP-A- 0583308 JP-T- 6506933	20-07-93 21-12-92 30-10-92 23-02-94 04-08-94
FR-A-2676361	20-11-92	NONE	
EP-A-373988	20-06-90	LU-A- 87399 CA-A- 2004309 JP-C- 1770249 JP-A- 3014509 JP-B- 4060569	10-07-90 02-06-90 30-06-93 23-01-91 28-09-92
WO-A-9208685	29-05-92	FR-A- 2669023 FR-A- 2680171 AT-T- 117665 CA-A- 2073353 DE-D- 69107056 DE-T- 69107056 EP-A- 0510165 ES-T- 2067255 US-A- 5362494	15-05-92 12-02-93 15-02-95 15-05-92 09-03-95 18-05-95 28-10-92 16-03-95 08-11-94

